

Correlated V/R and IR photometric variations in the Be/X-ray binary LS I +61° 235/RX J0146.9+6121

P. Reig^{1,2}, I. Negueruela³, M.J. Coe⁴, J. Fabregat⁵, A.E. Tarasov⁶ and R. K. Zamanov^{7,8}

¹ *Foundation for Research and Technology-Hellas. 711 10 Heraklion. Crete. Greece.*

² *Physics Department. University of Crete. 710 03 Heraklion. Crete. Greece.*

³ *SAX SDC, Agenzia Spaziale Italiana, c/o Telespazio, via Corcolle 19, 00131 Roma, Italy.*

⁴ *Physics & Astronomy Department, Southampton University, Southampton, SO17 1BJ, UK.*

⁵ *Departament d'Astronomia i Astrofísica, Universitat de València, E-46100 Burjassot-València, Spain.*

⁶ *Crimean Astrophysical Observatory, 334413 Nauchny, Crimea, Ukraine.*

⁷ *National Astronomical Observatory Rozhen, P.O. Box 136, BG-4700 Smoljan, Bulgaria.*

⁸ *Departamento de Física, Universidad de Jaén, C/Virgen de la Cabeza 2, 23071 Jaén, Spain.*

Received ????????; accepted ?????????

ABSTRACT

We report on the long-term variability of the Be/X-ray binary LS I +61° 235/RX J0146.9+6121. New optical spectroscopic and infrared photometric observations confirm the presence of global one-arm oscillations in the circumstellar disc of the Be star and allow us to derive a V/R quasi-period of 1240 ± 30 days. Pronounced shell events, reminiscent of the spectacular variations in Be stars, are also seen. We have found that the J , H and K infrared photometric bands vary in correlation with the spectroscopic V/R variations, implying that the one-armed disc oscillations are prograde. The effect of the oscillations is not only seen in the $H\alpha$ line but also in the He I $\lambda 6678$ and Paschen lines. Since these lines are formed at different radii in the equatorial disc of the Be star, such effect confirms the global nature of the perturbation. The Keplerian disc has been found to be denser than the average of a sample of isolated Be stars, which may be indicative of some kind of interaction with the compact companion. Finally, from a *Rossi X-ray Timing Explorer* observation we derive a spin period of the neutron star of 1404.5 ± 0.5 s.

Key words: stars: individual: LS I +61° 235 – binaries: general – X-rays: stars – stars: pulsars: individual: RX J0146.9+6121 – stars: emission-line, Be

1 INTRODUCTION

RX J0146.9+6121 is one of the slowest high mass X-ray pulsar systems (Mereghetti et al. 1993; Hellier 1994). Its optical counterpart is the $V = 11.2$, B1III-V star LS I +61° 235 located at an estimated distance of 2.3 ± 0.5 kpc (Motch et al. 1997; Coe et al. 1993; Reig et al. 1997a, hereafter R97). R97 derived the astrophysical parameters of the optical counterpart and reported V/R variations with a quasi-period of ~ 3 years. The line profile variability was attributed to the prograde precession of a one-armed mode confined in the Be star's disc.

Be/X-ray binaries comprise approximately 70% of the more general class of high mass X-ray binaries, the other $\sim 30\%$ containing evolved (luminosity class I and II) primaries. In a Be/X-ray binary the optical companion, a Be star, is characterised by an emission line spectrum and an infrared excess when compared to normal B-type stars of the same spectral type. These two observational properties

have their origin in the cool gaseous quasi-Keplerian disc which lies on the equatorial plane of the central star. A neutron star revolves around the $10\text{--}20\text{ M}_{\odot}$ primary in a rather eccentric orbit ($e \sim 0.2\text{--}0.8$) and accretes material expelled by the Be star from the disc – in the form of a low velocity, high density equatorial wind – giving rise to the X-rays.

The physical properties of the Be star's disc in Be/X-ray binaries have traditionally been considered to be the same as those of isolated Be stars. Indeed, the long-term variability characterised as disc-loss phases and V/R variations have been seen in both, isolated and Be/X-ray binaries (Okazaki 1997; Negueruela et al. 1998). However, it is not yet clear what role the compact companion in Be/X-ray binaries may play in the onset and subsequent development of the perturbation that gives rise to the asymmetric profiles or in the formation and loss of the disc. There is growing evidence that the circumstellar disc surrounding isolated Be stars and Be star in X-ray binaries may not share, on average,

Table 1. Journal of spectroscopic observations and spectral parameters for the H α line. Errors are $\lesssim 8\%$

Date	MJD	Tele scope	Range nm	EW Å	ΔV km/s	profile
17-08-94	49582	BNAO ^a	650-662	-8.4	252	V>R
08-06-96	50243	BNAO ^a	650-662	-9.6	257	V<R
09-06-96	50244	BNAO ^a	650-662	-10.1	258	V<R
01-08-96	50297	INT ^a	610-700	-10.1	316	V<R
26-08-96	50322	BNAO ^a	650-662	-10.2	298	V<R
13-11-96	50401	CAO	653-660	-10.3	259	V<R
12-12-96	50430	CAO	653-660	-10.3	251	V<R
01-02-97	50481	INT ^a	580-740	-11.9	288	V<R
21-08-97	50682	INT ^a	610-700	-12.1	242	V>R
18-09-97	50710	CAO	653-660	-12.0	223	V>R
10-10-97	50732	BNAO ^a	650-662	-10.7	233	V>R
28-10-97	50750	JKT	638-673	-11.8	258	V>R
10-02-98	50855	INT	640-680	-11.2	235	V>R
01-10-98	51088	BNAO ^b	650-670	-11.0	337	V>R
04-10-98	51091	BNAO ^b	650-670	-11.3	333	V>R
08-10-98	51095	WHT	649-661	-11.5	336	V>R
22-07-99	51382	CAO	653-660	-10.8	298	shell
25-07-99	51385	SKI	555-756	-11.9	312	shell
26-07-99	51386	SKI	555-756	-11.6	325	shell
19-08-99	51410	INT ^b	624-680	-11.2	298	shell
17-09-99	51439	BNAO ^b	650-670	-12.2	286	shell
19-09-99	51441	BNAO ^b	650-670	-11.7	285	shell
BNAO ^a : ISTA 580×400 + 632 l/mm				0.2 Å/pix		
BNAO ^b : Photometrics 1024×1024 + 632 l/mm				0.2 Å/pix		
CAO: Electronix 1024×260 CCD + 600 l/mm				0.06 Å/pix		
JKT: TEK4 + RBS + 2400 l/mm				0.4 Å/pix		
INT ^a : TEK3 + IDS + 235 mm + R1200 l/mm				0.8 Å/pix		
INT ^b : EEV10 + IDS + 500 mm + R1200 l/mm				0.2 Å/pix		
WHT: UES + 31 l/mm, R=54000				0.06 Å/pix		
SKI: ISA608 2000×800 + 1300 l/mm				1 Å/pix		

the same physical properties (Reig et al. 1997b; Negueruela et al. 1998; 1999)

In this paper we analyse optical, infrared and X-ray data and search for correlations between the characteristics of the radiation at these wavelengths. Our ultimate goal is to assess the validity of the global one-armed oscillation model and whether the compact companion has any effect on the V/R variability.

2 OBSERVATIONS AND RESULTS

2.1 Optical data

Optical spectroscopic observations were made from the 2.6m telescope at the Crimean Astronomical Observatory (CAO), the 2m RCC telescope of the Bulgarian National Astronomical Observatory "Rozhen" (BNAO), the 1.3m telescope of the Skinakas Observatory (SKI), in Crete (Greece), the 1.0m Jacobus Kapteyn Telescope (JKT), the 2.5m Isaac Newton Telescope (INT) in service mode and the 4.2m William Herschel Telescope (WHT). The last three telescope are located in the Observatorio del Roque de Los Muchachos (La Palma, Spain). Table 1 shows the journal of the observations and the instrument set-up. For a description of the observations not mentioned in the table see R97.

Fig 1 shows a selected sample of H α line profiles of LS

Table 2. Journal of infrared observations

Date	MJD	J	H	K
13-01-96	50096.42	9.86±0.03	9.58±0.02	9.35±0.02
19-03-96	50162.36	9.78±0.02	9.60±0.01	9.40±0.02
28-07-96	50293.68	9.73±0.05	9.51±0.06	9.34±0.04
29-07-96	50294.65	9.73±0.10	9.51±0.09	9.31±0.08
18-07-97	50648.65	9.83±0.02	9.56±0.02	9.32±0.02
19-07-97	50649.66	9.77±0.01	9.52±0.01	9.32±0.01
21-07-97	50651.66	9.76±0.02	9.53±0.02	9.29±0.02
17-06-98	50982.72	10.01±0.01	9.81±0.02	9.53±0.02
27-10-98	51114.42	10.01±0.02	9.69±0.02	9.48±0.02
28-10-98	51115.67	9.82±0.02	9.66±0.02	9.45±0.02
28-10-98	51115.67	9.85±0.02	9.65±0.02	9.44±0.02
26-07-99	51386.72	9.80±0.02	9.58±0.02	9.37±0.02
27-07-99	51387.66	9.96±0.03	9.67±0.02	9.43±0.02
29-07-99	51389.63	9.85±0.02	9.61±0.02	9.36±0.02
30-07-99	51390.66	9.81±0.02	9.58±0.02	9.34±0.02
01-08-99	51392.64	9.90±0.04	9.63±0.03	9.41±0.02
02-10-99	51454.49	9.79±0.02	9.54±0.02	9.33±0.02
03-10-99	51455.60	9.86±0.02	9.58±0.02	9.32±0.02
04-10-99	51456.56	9.82±0.02	9.57±0.02	9.31±0.02

I +61° 235 covering the period 1991 August – 1999 September. Part of the new data (all observations taken after 1996 February plus those from Rozhen BNAO) are plotted together with some of the spectra presented in R97. The evolution of the V/R ratio can be seen in Fig 2a. Different shapes of the line have been represented by different symbols. Stars are used for $V > R$ points, triangles for $V < R$ phases, dots for single-peak lines and squares for shell profiles (i.e. when the central absorption goes beyond the continuum). Note that the shell phase is very brief and always occurs during the transition from $V > R$ to $V < R$. In contrast, the transition from $V < R$ to $V > R$ is separated by single-peak profiles. Strictly speaking there are not symmetric single-peak lines in our observations (as observed in pole-on stars) since one can always see flank inflections revealing a second peak. Nevertheless, we will refer to as single-peak phase the profiles seen during the transition $V < R$ to $V > R$. By fitting a sine function to the V/R curve we refined the V/R quasi-period to a value of 1240 ± 30 days.

We have also measured the separation of the blue and red peaks by fitting two Gaussian functions to the line profile (Fig 2d). The peak separation gives a measure of the velocity field, assuming it to be Keplerian. The blue-dominated profiles seem to sample a wider range of velocities than red-dominated profiles: $V > R$ points spread over a velocity range 220–360 km s⁻¹, whereas $V < R$ points distribute around 250–320 km s⁻¹. Also, blue-dominated profiles with high values of the peak separation occur only at the end of the $V > R$ cycle, just before the shell phase.

The density wave does not only affect the H α line but also the Paschen series and He I $\lambda 6678$ (Fig 3). Since these lines are formed at different regions inside the equatorial disc the perturbation must extend over a very wide region, hence confirming its global nature.

Figure 1. Evolution of the H α line profile over the period 1991 August 28 – 1999 September 17 (MJD 48497–51439)

2.2 Infrared data

The infrared observations were taken with the Continuous Variable Filter (CVF) infrared photometer, using the 1.5-m Carlos Sánchez telescope, located at the Teide Observatory, in Tenerife (Spain). The data were reduced following the procedure described by Manfroid (1993). Instrumental values were transformed to the TCS standard system (Alonso et al. 1998). Table 2 shows the results of the infrared observations. For earlier observations the reader is referred to R97.

The combination of the old and new infrared data yields a very interesting result, namely the correlation between the V/R variations and the infrared magnitudes. Fig 2 shows the J and K light curves and the evolution of the colour index J-K, covering the period 1991 August to 1999 October. The light curves were rebinned into 30 day bins. The errors were calculated from the photometric errors depending on the number of points, N , in each bin: when the corresponding bin contained only one point we simply took the observation error, if $1 < N \leq 5$ then we defined the error as $\sum |x_i - x_m|/N$ and if $N > 5$ then we considered the standard deviation $\sqrt{(x_i - x_m)^2/N}$.

As can be seen in Fig 2, there is a distinct modulation with an amplitude, from maximum to minimum, of ~ 0.3 mag. The period of this modulation is 3.1–3.8 years, in good agreement with the optical V/R variability of 3.4 years. Interestingly, the infrared maxima occur in coincidence with the optical ($\sim$ MJD 49250) and X-ray ($\sim$ MJD

50630) outbursts. On the other hand, the infrared colours do not change drastically over the period of the observations. That is, the slope of the infrared continuum remained the same over the time covered by the observations.

2.3 X-ray data

LS I +61° 235 is a persistent low-luminosity Be/X-ray binary. These systems are characterised by long pulse periods, low X-ray variability ($L_{max}/L_{min} \lesssim 10$) and low but permanent levels of X-ray emission (Reig & Roche 1999).

RX J0146.9+6121 was observed with the RXTE *Proportional Counter Array* (PCA) on March 21, 1998 (02:01–11:54 UT). Good time intervals were defined by removing data taken at low Earth elevation angle ($< 10^\circ$) and during times of high particle background. An offset of only 0.02° between the source position and the pointing of the satellite was allowed, to ensure that any possible short stretch of slew data at the beginning and/or end of the observation was removed. All five PCA units were functioning during the entire observation. The total net exposure was 19223 s. Due to the relatively wide field of view of the PCA instrument (1° FWHM) the nearby X-ray source 4U 0142+61 (White et al. 1996) also contributed to the total flux. After correcting for collimator efficiency the contribution of 4U 0142+61 to the total observed flux in the energy range 2–30 keV was estimated to be of $\sim 10\%$. Because of this no X-ray spectral analysis was attempted.

The pulse period was determined by correcting the data

Figure 2. Evolution of the V/R ratio (a), J band (b), K band (c), J-K index (d), H α equivalent width (e) and peak separation (f). A correlation between the V/R cycle and the infrared emission is apparent. The arrow in panel (e) indicates the X-ray outburst

to the solar system barycentre and using the epoch folding technique, i.e., we folded the data over a range of periods and searching for a maximum χ^2 as a function of period. The pulse period found was 1404.5 ± 0.5 s, which is virtually the same as the 1404.2 ± 1.2 s pulse period obtained in another RXTE observation nine months earlier (Haberl et al. 1998).

3 DISCUSSION

3.1 Global $m = 1$ oscillations

In R97 we investigated the different models that had been put forward to explain the V/R variability in Be stars and

Figure 3. The V/R oscillations are not only seen in H α (left) but also in He I $\lambda 6678$ (middle) and Paschen lines Pa11 ($\lambda 8863$) and Pa12 ($\lambda 8750$) (right)

concluded that the model which best accounted for the observational data in LS I +61° 235 was the Global One-armed Oscillation model (Okazaki 1991, 1997; Papaloizou et al. 1992). This model suggests that the long-term V/R variations are caused by global $m = 1$ oscillations in the cool equatorial disc of the Be star. In other words, an enhanced density perturbation develops on one side of the disc, which slowly precesses. The precession time being that associated with the V/R quasi-period. The density perturbation is confined within a few stellar radii in the disc and the precession period turns out to be fairly insensitive to the size of the disc (Savonije & Heemskerk 1993).

One prediction of the model is that no changes in the slope of the infrared continuum are expected. The reason is that the V/R variations are not the result of changes in the radial gradient of the circumstellar gas. The slope of the infrared continuum is a measure of the radial density distribution but since matter in the disc does not move radially no changes in the shape of the infrared continuum are expected. This is exactly the behaviour that we find in the case of LS I +61° 235. While the individual infrared photometric bands changed ($\Delta J \approx \Delta H \approx \Delta K \sim 0.3$ magnitudes) the infrared colours remained unchanged (Fig. 2).

In principle, the issue of whether the motion of the perturbation occurs in the same sense (prograde rotation) or opposite sense (retrograde rotation) to the stellar rotation can be found out from the observations. Telting et al. (1994) realised that a prograde revolution implies that the $V > R$ phase must be followed by a shell profile and a similar profile but with much less pronounced absorption feature (or possibly a single peak line if the inclination is low) during the transition from $V < R$ to $V > R$. These characteristic line shapes must translate into noticeable photometric variations. According to Mennickent et al. (1997), we should expect a minimum of brightness when $V = R$ prior to the $V < R$ phase if the motion is prograde and $V > R$ after $V = R$ if retrograde. In LS I+61 235 the minimum of brightness in the infrared photometric bands occurred dur-

ing the shell phase ($V = R$) before the $V < R$ phase began, confirming the prograde nature of the precession inside the disc.

However, models of one-armed global density waves cannot reproduce strong shell-non-shell transitions like the ones seen in LS I +61° 235 which are reminiscent of the so-called *spectacular variations* (Doazan et al. 1983). Such shell events seem to be rare a phenomenon which have been reported for only three Be stars: γ Cas, 59 Cyg and Pleione (Hummel 1998 and references therein), all of which are either binaries or suspected binaries. One possible explanation might be a *thick* disc in the region where the perturbation lies. When the perturbation is behind the star and for the right inclination angle, no shell event is seen because the disc in between the central star and the observer is thin and does not occult the star nor the perturbation. The shell phase would occur when the perturbation is at inferior conjunction since the thicker disc would hide the central star from the observer. An alternative explanation is given by Hummel (1998) who suggested a tilted or warped circumstellar disc with precessing nodal line in addition to the density wave (see also Porter 1998).

3.2 A dense circumstellar disc in LS I +61° 235

The works by Dachs et al. (1986) and Hanuschik et al. (1988) have shown that the equivalent width of $H\alpha$ line emission for Be stars increases with the effective disc radius. Since for rotationally dominated profiles $\Delta V/(2v \sin i)$ can be regarded as a measure of the radius of the $H\alpha$ emitting region (Huang 1972), we expect a correlation between the peak separation and the $H\alpha$ equivalent width. Hanuschik et al. (1988) derived the following law

$$\log \left(\frac{\Delta V}{2v \sin i} \right) = a \log(EW(H\alpha)) + b \quad (1)$$

where $v \sin i$ is the projected rotational velocity and $EW(H\alpha)$ is given in angstroms. a and b are related to the rotational law index $a = -j/2$ ($j=0.5$ for Keplerian rotation and $j=1$ for conservation of angular momentum) and with the disc electron density, respectively. A least square fit to the LS I +61° 235 data gave $a = -0.23 \pm 0.10$, i.e. $j \approx 0.5$ and $b = +0.1 \pm 0.1$. These values are to be compared with the average values $a = -0.4$ and $b = -0.1$ found by Hanuschik et al. (1988) for a sample of 26 isolated Be stars. The higher value of b in LS I +61° 235 implies a denser disc than those of isolated Be stars.

The main (and probably only) difference between a Be/X-ray binary and an early-type isolated Be star is the presence of a neutron star in the former. It then seems natural to attribute the dissimilarity in the properties of the circumstellar envelopes to the influence of such compact companion. The neutron star trims the disc to a certain radius and prevents its free growth, making it denser. Disc truncation has been suggested in other Be/X-ray binaries like V0332+53 (Negueruela et al. 1999). This idea would support the hypothesis that the neutron star plays a fundamental role in the evolution and properties of the equatorial disc in Be/X-ray binaries, as proposed by Reig et al. (1997b).

3.3 X-ray/optical/IR correlations

In R97 the observation of an optical outburst around 1993 September-October (MJD ~ 49260) was reported. During the outburst the $H\alpha$ equivalent width, $EW(H\alpha)$, changed by $\sim 10 \text{ \AA}$, in about 270 days, decreasing to pre-outburst values ($\sim -8 \text{ \AA}$) in about the same amount of time (Fig 2c). This increased coincided with a single-peak phase of the $H\alpha$ profile. The question of whether this outburst was an isolated event or was associated with the V/R cycle remained open due to the short coverage of the data. The new observations show no new $EW(H\alpha)$ maximum. After the outburst the $EW(H\alpha)$ increased slowly up to a level of $-12 \text{ \AA}$, considerably lower than the peak of 1993 September. The new single-peak phase should have occurred during 1997 March-May. Unfortunately, the star was too close to the Sun to be observed. However, we notice that $EW(H\alpha)$ seems to have reached a maximum value just before and after the period when we would expect the single-peak phase (around MJD 50500). Thus, we are inclined to think that the higher $EW(H\alpha)$ associated with single-peak profiles reported in R97 may reflect the fact that the fluxes of both components are adding up and are not affected by the absorption feature present in the other profiles. In this context then, the increase in $EW(H\alpha)$ is real and would be an event related to the motion of the density pattern in the disc, rather than an isolated episode.

In 1997 July (MJD 50634) LS I +61° 235 underwent a small X-ray outburst (Haberl et al. 1998). The X-ray luminosity increased by nearly a factor of five in one week reaching $3.45 \times 10^{35} \text{ erg s}^{-1}$ in the energy range 0.5–10 keV. This outburst is marked with an arrow in Fig 2c. Interestingly, the outburst occurred at the time of the expected infrared and $EW(H\alpha)$ maxima. One is then tempted to attribute this correlated X-ray/optical/IR behaviour to the high density perturbation where most of the Balmer emission is formed. If the inclination of the system is less than 90° , when the high-density part of the equatorial disc is behind the star it offers the largest geometric area (especially so if the disc is thicker in this region) and the highest optical and infrared emission. If the neutron star happens to be close to the Be star it will accrete from this high-density material and the X-ray emission will be enhanced. New optical and X-ray observations around the next expected maximum are needed to solve this issue.

4 CONCLUSION

Optical spectroscopic observations confirm the presence of global one-armed oscillations in the circumstellar disc of LS I +61° 235. These oscillations manifest themselves as quasi-periodic variations in the shape of the $H\alpha$ line, whose asymmetric double peak profile alternates between red- ($V < R$) and blue-dominated ($V > R$) emission. The system also goes through Be-Be shell transitions, which might indicate an asymmetric vertical structure of the disc in the form of a thick or a tilted disc. The V/R quasi-period is determined to be $\sim 1240 \pm 30$ days. We have found a correlation between the infrared emission and the V/R variations. This is the first time that such correlation is reported in a Be/X-ray binary. From the pattern traced by the IR light curves in relation to the V/R ratio we conclude that the one-armed disk

oscillations are prograde. The Be star's disc in LS I +61° 235 is found to be denser than isolated Be stars, which may be connected to the presence of the neutron star. From a *Rossi X-ray Timing Explorer* observation we derive a spin period of the neutron star of 1404.5 ± 0.5 s.

Acknowledgements

We thank Chris Moran for providing us with one of the INT spectra and Dr E. V. Paleologou for helping us with the spectroscopic observations at Skinakas Observatory. Skinakas Observatory is a collaborative project of the University of Crete, the Foundation for Research and Technology-Hellas and the Max-Planck-Institut für Extraterrestrische Physik. P. Reig acknowledges support via the European Union Training and Mobility of Researchers Network Grant ERBFMRX/CT98/0195. IN is supported by an ESA external fellowship. R. Zamanov acknowledges support from Dirección general de relaciones culturales y científicas, Spain. Some observations were taken as part of the ING service observing programme. We are grateful to the referees, Dr D. Baade and Dr T. Rivinius, for useful comments.

REFERENCES

- Alonso A., Arribas S., Martínez-Roger C. 1998, A&AS 131, 209
 Coe M.J., Everall C., Norton A.J., Roche P., Unger S.J., Fabregat J., Reglero V. & Grunsfeld J.M., 1993, MNRAS, 261, 599
 Dachs, J., Hanuschik, R., Kaiser, D. & Rohe, D. 1986, A&A, 159, 276
 Doazan V., Franco M., Rusconi L., Sedmark G., Stalio R., 1983, A&A, 128, 171
 Haberl F., Angelini L. & Motch C., 1998, A&A, 335, 587
 Hanuschik R.W., Kozok J.R., Kaiser D., 1988, A&A 189, 147
 Hellier C., 1994, MNRAS, 271, L21
 Huang, S. 1972, ApJ, 171, 549
 Hummel W., 1998, A&A 330, 243
 Manfroid J. 1993, A&A 271, 714
 Mennickent, R. E., Sterken, C. & Vogt, N. 1997, A&A, 326, 1167
 Mereghetti S., Stella L. & De Nile F., 1993, A&A, 278, L23
 Motch C., Haberl F., Dennerl K., Pakull M. & Janot-Pacheco E., 1997, A&A, 323, 853
 Negueruela I., Reig P., Coe M.J. & Fabregat J., 1998, A&A, 336, 251
 Negueruela I., Roche P., Fabregat J., Coe, M. J., 1999, MNRAS, 307, 695
 Okazaki A.T., 1991, PASJ 43, 75
 Okazaki A.T., 1997, A&A 318, 548
 Papaloizou, J.C., Savonije, G. J., Henrichs, H. F, 1992, A&A, 265, L45
 Porter J.M., 1998, A&A 336, 966
 Reig P., Fabregat J., Coe M. J., Roche P., Chakrabarty D., Negueruela I & Steele I., 1997a, A&A, 322, 183, (R97).
 Reig P., Fabregat J. & Coe M.J., 1997b, A&A, 322, 193
 Reig P. & Roche P., 1999, MNRAS, 306, 100
 Savonije G.J. & Heemskerk M.H.M., 1993, A&A, 276, 409
 Telting, J.H., Heemskerk M.H.M., Henrichs H.F. & Savonije G.J., 1994, A&A, 288, 558
 White N.E., Angelini, L., Ebisawa K., Tanaka Y. & Ghosh P., 1996, ApJ, 463, L83





